



A characteristic language for rational w-power

Sandrine Julia

► To cite this version:

Sandrine Julia. A characteristic language for rational w-power. Developments in Language Theory, Jul 1997, Greece. p. 299-308, 1997. <hal-00258666>

HAL Id: hal-00258666

<https://hal.archives-ouvertes.fr/hal-00258666>

Submitted on 23 Feb 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A characteristic language for rational ω -power

Sandrine JULIA

Laboratoire I3S-U.R.A. 1376-C.N.R.S.,
Université de Nice - Sophia Antipolis,
Route des Colles, B.P. 145
06903 SOPHIA ANTIPOLIS Cedex
France
e-mail: `julia@diamant.unice.fr`

Abstract

Consider a rational ω -power L^ω over a finite alphabet Σ and call $\chi(L^\omega)$ the following language: $\chi(L^\omega) = \{u \in \Sigma^+, uL^\omega \subseteq L^\omega \text{ and } u^\omega \in L^\omega\}$. It is already known that this language is rational and an upper bound of the set of ω -generators of L^ω , itself ω -generator of L^ω as soon as it is a semigroup.

We prove that two rational ω -powers are equal if and only if they have the same language χ . We deduce that two ω -languages, both ω -power and adherence are equal if and only if they have the same set Per_χ where Per_χ denotes the set of periodic ω -words with a period in χ . At last, we present some additionnal results especially in the case of rational deterministic ω -power.

Introduction

A rational ω -language is a set of right infinite words which can be recognized by a Büchi automaton. Actually, the class of rational ω -languages is obtained as a closure of the class of finite ω -languages and Büchi proved that it is also the class of ω -languages recognized by finite Büchi automata [2, 4]. However, a rational ω -language is not necessarily recognized by a deterministic Büchi automaton. In spite of the existence of Muller automata which ensure that every rational ω -language is recognized by a deterministic automaton, there is no way to define a canonical minimal automaton. Moreover, the syntactic congruence of a rational ω -language is finite but the converse does not hold and two distinct ω -languages can have the same syntactic congruence [1].

The rational ω -languages are of the form $\bigcup_{i=1}^n A_i B_i^\omega$, with A_i and B_i some rational languages and so we intend to investigate this operation $^\omega$ which is a generalization of the Kleene's star operation. In this way, the following consequence of Büchi theorem provides us some interesting results. A rational ω -language is entirely characterized by its set of periodic ω -words. Hence, we are able to prove that a rational ω -power, that is an ω -language of the form L^ω for some language L , is entirely characterized by the finitary language $\chi(L^\omega) = \{u \in \Sigma^+, uL^\omega \subseteq L^\omega \text{ and } u^\omega \in L^\omega\}$. This language was previously known as an upper bound of the set of the ω -generator of a rational ω -language, itself rational if the

ω -language is rational. Moreover, it is an ω -generator if it is a semigroup. In particular, it is true if the considered ω -language is an adherence [6]. We derive from this some results about rational ω -languages which are both ω -powers and adherences.

Another consequence deals with rational deterministic ω -languages. The set of ω -generators of a rational ω -language admits a finite number of maximal ω -generators [7]. In addition, we prove that in the case of deterministic ω -languages, the intersection of those maximal ω -generators is still an ω -generator.

1 Definitions and notation

Let Σ be a finite alphabet. A *word* (resp. ω -*word*) is a finite (resp. infinite) concatenation of letters in Σ . We note ε the empty word. Σ^* is the set of words over Σ , $\Sigma^+ = \Sigma^* \setminus \{\varepsilon\}$. Σ^ω is the set of ω -words. The subsets of Σ^* are called *languages*, those of Σ^ω *ω -languages*. Let L be a language, the language L^* is the set of words built with words in L : $L^* = \{\varepsilon\} \cup \{a_1 \dots a_n / \forall i \ 1 \leq i \leq n, a_i \in L\}$. In the same way, the ω -*power* L^ω is the set of ω -words: $L^\omega = \{a_1 \dots a_n \dots / \forall i > 0, a_i \in L \setminus \{\varepsilon\}\}$. L^* (resp. L^ω) is generated (resp. ω -generated) by L , and so L is called a *generator* (resp. ω -*generator*).

A word u is *prefix* of v if $v \in u(\Sigma^* \cup \Sigma^\omega)$ and we write: $u < v$. The induced order is the *prefix order*. For $v \in (\Sigma^* \cup \Sigma^\omega)$, $\text{Pref}(v)$ stands for the set of all prefixes of v . Hence, for every $L \subseteq (\Sigma^* \cup \Sigma^\omega)$, $\text{Pref}(L)$ is the set of the prefixes of the words in L . An ω -word $a_1 \dots a_n \dots$ is *periodic* (resp. *ultimately periodic*) if $\exists p \geq 1$ such that $\forall i > 0 \ a_i = a_{i+p}$ (resp. $\exists p \geq 1$ such that $\exists m \geq 1 \ \forall i \geq m \ a_i = a_{i+p}$).

Let L be a language, the *stabilizer* of L^ω is the following language: $\text{Stab}(L^\omega) = \{u \in \Sigma^*, uL^\omega \subseteq L^\omega\}$. $\text{Stab}(L^\omega)$ is a submonoid which can be understood as the set of all prefixes of ω -words in L^ω that could be fixed to the ω -words of L^ω to obtain yet another element in L^ω . Moreover every ω -generator of L^ω is included in $\text{Stab}(L^\omega)$ and so, when $\text{Stab}(L^\omega)$ is an ω -generator of L^ω , it is the greatest [7].

Let $L \subseteq \Sigma^*$. An *L -factorization* of a word u in L^+ is a finite sequence of words in $L \setminus \{\varepsilon\}$: (u_1, u_2, \dots, u_n) such that $u = u_1 u_2 \dots u_n$. An *L -factorization* of an ω -word w in L^ω is an infinite sequence of words in $L \setminus \{\varepsilon\}$: $(w_1, w_2, \dots, w_n, \dots)$ such that $w = w_1 w_2 \dots w_n \dots$. We will say indifferently *L -factorization* or *factorization over L* .

Let $L \subseteq \Sigma^*$. An *L -factorization* (u_1, u_2, \dots) of an ω -word is said to be *periodic* (resp. *ultimately periodic*) if there exists $p > 0$ such that $\forall i > 0 \ u_i = u_{i+p}$ (resp. $\exists j > 0 \ \forall i \geq j \ u_i = u_{i+p}$).

Let L be a language, the *adherence* of L is the ω -language $\text{adh}(L) = \{w \in \Sigma^\omega, \text{Pref}(w) \subseteq \text{Pref}(L)\}$ [9].

Let L be an ω -language, $\text{Per}(L)$ denotes the set of periodic ω -words in L whereas $\text{Ult}(L)$ is the set of ultimately periodic ω -words in L .

A *Büchi automaton* \mathcal{A} is a finite automaton $(\Sigma, Q, \delta, I, T)$ where Σ denotes the finite alphabet, Q the finite set of states, $I \subseteq Q$ the set of initial states and $T \subseteq Q$ the set of recognition states. A *run* of an ω -word w in \mathcal{A} is an infinite sequence $l = (q_i)_{i \geq 0}$ of elements in Q such that $\delta(q_i, w_{i+1}) = q_{i+1}$, with w_i the i^{th} letter of w . An ω -word w belongs to the ω -language recognized by \mathcal{A} if there exists a run $(q_i)_{i \geq 0}$ such that $q_0 \in I$ and $\text{Inf}(w) \cap T \neq \emptyset$, where $\text{Inf}(w) = \{q, \text{Card}(\{i / q_i = q\}) \text{ is infinite}\}$. More intuitively, an ω -word w is accepted if its run in \mathcal{A} enters infinitely often in a recognition state.

2 Useful results

We recall that rational ω -languages are the ω -languages of the form: $L = \bigcup_{i=1}^n A_i B_i^\omega$ with $n \geq 1$ such that for every i , A_i and B_i are rational languages, subsets of Σ^+ . This is also the class of ω -languages recognized by Büchi automata, as it was proved by Büchi [2], as a Kleene's theorem for the infinitary case. Moreover, this set is a boolean algebra [8] and so, there is the following consequence of Büchi's theorem:

Proposition 1 [2]

- (1) Let L be a rational ω -language. If $L \neq \emptyset$ then $Ult(L) \neq \emptyset$.
- (2) Let L and M be two rational ω -languages. Then $L = M$ if and only if $Ult(L) = Ult(M)$.

Proof (1) If L is a rational ω -language, then $L = \bigcup_{i=1}^n A_i B_i^\omega$ for some rational languages A_i, B_i in Σ^+ . If $L \neq \emptyset$, then we can assume that there exist non-empty words $x \in A_1$ and $y \in B_1$, such that $xy^\omega \in L$.

(2) Let L and M be two rational ω -languages such that $Ult(L) = Ult(M)$. The symmetrical difference $L \Delta M = (L \cup M) \setminus (L \cap M)$ is a rational ω -language because the set of rational ω -languages is a boolean algebra. Furthermore, $L \Delta M$ does not contain any ultimately periodic ω -word. Furthermore, a non-empty rational ω -language necessarily contains an ultimately periodic ω -word, as shown above. Thus, we have $L \Delta M = \emptyset$ and thus $L = M$. \square

Below, we recall some result concerning adherences we need in the sequel.

Corollary 1 [6] Let L and M two rational adherences. $L = M$ if and only if $Pref(L) = Pref(M)$.

3 The language χ

Let L^ω be a rational ω -power, we denote by χ the following language:

$$\chi(L^\omega) = \{u \in \Sigma, uL^\omega \subseteq L^\omega \text{ and } u^\omega \in L^\omega\}$$

This language is already known as an upper bound of the set of the ω -generators of a rational ω -power. Moreover, it is an ω -generator as soon as it is a semigroup [6]. In the sequel, we prove that it is characteristic of such a rational ω -power.

We present some notation, depending on the language χ . If L^ω is an ω -power, $Per_\chi(L^\omega)$ denotes the set of its periodic ω -words which have a period in $\chi(L^\omega)$. Recall that $Ult(L^\omega)$ denotes the set of its ultimately periodic ω -words. Remark that $Per_\chi(L^\omega) = \{u^\omega, u \in \chi(L^\omega)\}$ but is not necessarily the set of *all* periodic ω -words of L^ω . Equivalently, $Per_\chi(L^\omega) = \{w \in \Sigma^\omega, \exists u \in \Sigma^+ \quad uL^\omega \subseteq L^\omega \text{ and } w = u^\omega\}$.

Example Let $L = a + ba$. $\chi(L^\omega) = (a + ba)^+$, $Per_\chi(L^\omega) = \{u^\omega, u \in (a + ba)^+\}$ while the periodic ω -word $(ab)^\omega \in Per(L^\omega) \setminus Per_\chi(L^\omega)$.

Another consequence of Büchi's theorem is about the rationality of the set of ultimately periodic ω -words of a rational ω -power.

Lemma 1 [2] *Let L be a rational ω -language such that $L \neq \text{Ult}(L)$. Then $\text{Ult}(L)$ is not a rational ω -language.*

Proof If $\text{Ult}(L)$ were a rational ω -language, then the equality $\text{Ult}(L) = \text{Ult}(\text{Ult}(L))$ would imply the equality $L = \text{Ult}(L)$. \square

Lemma 2 [3] *Let L be a language. Every ultimately periodic ω -word in L^ω has an ultimately periodic L -factorization.*

Proof Let $w = uv^\omega \in L^\omega$. The L -factorization of uv^ω infinitely often “cuts” the word $v = v_1 \dots v_n$, $v_i \in \Sigma$. Nevertheless, v being of finite length, then it cannot be cut in more than in a finite number of different places. In fact, there exist i, j and k three non-negative integers such that $w_1 \dots w_i \in L^*$ with $w_1 \dots w_i = uv_1 \dots v_{k-1}$, $w_{i+1} \dots w_j \in L^*$ with $w_{i+1} \dots w_j \in \{v_k \dots v_n v^i v_1 \dots v_{k-1}, i \geq 0\}$ and $w = w_1 \dots w_i (w_{i+1} \dots w_j)^\omega$. We thus obtain an ultimately periodic L -factorization of L^ω . \square

Now we can propose our main result:

Theorem 1 *Let L and M be two rational languages. The following assertions are equivalent:*

- (i) $L^\omega = M^\omega$
- (ii) $\chi(L^\omega) = \chi(M^\omega)$
- (iii) $\text{Stab}(L^\omega) = \text{Stab}(M^\omega)$ and $\text{Per}_\chi(L^\omega) = \text{Per}_\chi(M^\omega)$

Proof It is clear that (i) \Rightarrow (ii) and (i) \Rightarrow (iii). (ii) \Rightarrow (i): we know that a rational ω -language is characterized by its ultimately periodic ω -words. Let $w = uv^\omega$ be an ω -word in L^ω . From lemma 2, w has an ultimately periodic L -factorization thus there exist $u' \in L^*$ and $v' \in L^+$ such that $w = u'v'^\omega$. $L^* \subseteq \chi(L^\omega) = \chi(M^\omega)$. We deduce that $u'M^\omega \subseteq M^\omega$ and $v'^\omega \in M^\omega$ which imply that $w \in M^\omega$. Thus, we have just proved that $\text{Ult}(L^\omega) \subseteq \text{Ult}(M^\omega)$. In the same way, one can prove that $\text{Ult}(M^\omega) \subseteq \text{Ult}(L^\omega)$. Hence we get the equality $L^\omega = M^\omega$. (iii) \Rightarrow (i): if $\text{Stab}(L^\omega) = \text{Stab}(M^\omega)$ and $\text{Per}_\chi(L^\omega) = \text{Per}_\chi(M^\omega)$, one can apply the previous argument once more. \square

Observe that the equality between the stabilizers is not sufficient to ensure the equality between two rational ω -powers [6], nor the equality of the sets of periodic ω -words, as the following example attests.

Example

- $L = a^*b$ and $M = a + b$: L^ω and M^ω have the same stabilizer equal to Σ^* but $a^\omega \in M^\omega \setminus L^\omega$. In addition, $a^\omega \in (\text{Per}_\chi(M^\omega) \setminus \text{Per}_\chi(L^\omega))$.
- $L = bc + a + (c^+a^*)^*b$ and $M = a + (c^+a^*)^*b$: their sets of periodic words are identical i.e. $\text{Per}(L^\omega) = \text{Per}(M^\omega) = \{u^\omega, u \in L^+\}$. Nevertheless, the word $bca^\omega \in L^\omega \setminus M^\omega$ and thus the two ω -languages are different. There, $\text{Per}_\chi(L^\omega) \neq \text{Per}_\chi(M^\omega)$ because $(bca)^\omega$ belongs to the first but not to the second.

An automaton which recognizes $\chi(L^\omega)$ can be built from the Büchi automaton which recognizes L^ω . Consequently, the previous result provides a new algorithm to decide the equality between two rational ω -powers.

The notion we use are still meaningful for ω -languages which are not ω -powers. Thus, the stabilizer of the ω -language K is defined as follows: $Stab(K) = \{u \in \Sigma^+, uK \subseteq K\}$ and $\chi(K) = \{u \in \Sigma^+, uK \subseteq K \text{ and } u^\omega \in K\} = \{u \in Stab(K), u^\omega \in K\}$. Notice that the language $\chi(K)$ is no longer characteristic of ω -languages K which are not ω -powers, as it can be seen in this example:

Example $K = a^+b^\omega + a^\omega$ and $K' = a^\omega$. But $\chi(K) = \chi(K')$.

Below we propose a description of $Ult(L^\omega)$, the set of ultimately periodic ω -words in L^ω :

Proposition 2 *Let L be a rational language. The following equalities hold:*

$$Ult(L^\omega) = \chi(L^\omega)Per_\chi(L^\omega) = Stab(L^\omega)Per_\chi(L^\omega)$$

Proof If $w \in Ult(L^\omega)$, $w = uv^\omega$, with $u \in L^*$, $v \in L^+ \subseteq \chi(L^\omega) \subseteq Stab(L^\omega)$, according to lemma 2. Thus, $v^\omega \in Per_\chi(L^\omega)$ and if $u \neq \varepsilon$, $u \in Stab(L^\omega)$. Conversely, we obtain the result simply from the definitions of $\chi(L^\omega)$, $Stab(L^\omega)$ and $Per_\chi(L^\omega)$. \square

Corollary 2 *Let L be a rational language such that $L^\omega \neq Ult(L^\omega)$, $Per_\chi(L^\omega)$ is not a rational ω -language.*

Proof Consider a rational language L . From proposition 2, if $Per_\chi(L^\omega)$ were rational, this would imply the rationality of $Ult(L^\omega)$ since the languages $\chi(L^\omega)$ and $Stab(L^\omega)$ are rational. This is wrong, as we already have observed in lemma 1. \square

More generally, if G is a rational ω -generator of L^ω and if $L^\omega \neq Ult(L^\omega)$, the set $Per_G(L^\omega)$ is not rational. Indeed, $Ult(L^\omega) = G^*Per_G(L^\omega)$ and the rationality of $Per_G(L^\omega)$ would imply the rationality of $Ult(L^\omega)$.

We previously knew that $Pref(L^\omega) = Pref(L^*) = Pref(\chi(L^\omega)) = Pref(Stab(L^\omega))$. We add:

Corollary 3 *Let L be a rational language, then*

$$Pref(\chi(L^\omega)) = Pref(Ult(L^\omega)) = Pref(Per_\chi(L^\omega))$$

Proof On one hand, from proposition 2, $Pref(\chi(L^\omega)) \subseteq Pref(Per_\chi(L^\omega)) \subseteq Pref(Ult(L^\omega)) \subseteq Pref(L^\omega)$. Conversely, $L^* \subseteq \chi(L^\omega)$ then $Pref(L^\omega) = Pref(L^*) \subseteq Pref(\chi(L^\omega))$ and we get the equalities. \square

4 Some consequences

4.1 Adherences

The next results describe some ways to characterize an ω -language which is simultaneously an adherence and an ω -power:

Proposition 3 [5, 6] *Let L be a language in Σ^* . The following assertions are equivalent:*

- (i) L^ω is an adherence
- (ii) $L^\omega = adh(L^*)$
- (iii) $adh(L) \subseteq L^\omega$

When L^ω is an adherence, it is well known that $Stab(L^\omega)$ and $\chi(L^\omega)$ are equal [6], and both of them are ω -generators. It is particularly the case when L^ω admits a finite ω -generator. Moreover, let L be a language in Σ^* , then L^ω is an adherence if and only if

$$adh(\chi(L^\omega)) = L^\omega$$

If L^ω is not an adherence, all of its ω -generators share the same adherence, namely $adh(\chi(L^\omega))$. In that case, this adherence does not characterize the ω -power. Indeed, it suffices for any two ω -powers to share the same set of prefixes to have the same adherence.

Example Let $L = ba^*$. Then $Stab(L^\omega) = \chi(L^\omega) = b\Sigma^*$ and $adh(Stab(L^\omega)) = b\Sigma^\omega \neq L^\omega$. Indeed, $ba^\omega \in (b\Sigma^\omega \setminus (ba^*)^\omega)$. Therefore L^ω is not an adherence ($ba^\omega \in adh(L)$ but $ba^\omega \notin L^\omega$). Let $M = a + b$ and $N = a^*b$. $M^\omega = adh(M^*) = adh(N^*) \neq N^\omega$.

Observe that the previous characterizations are not true anymore if we consider ω -languages which are not ω -powers as we can see in the example below.

Example Let $K = a^*b^\omega + a^\omega$. K is not an ω -power although it is an adherence. In addition, $Stab(K) = a^+ = \chi(L^\omega)$ and $adh(Stab(K)) = a^\omega$ which is different from K .

We will now prove that, in the same way a rational ω -power can be characterized by the language $\chi(L^\omega)$, a rational adherence of the form L^ω is characterized by the set of its periodic ω -words with a period in $\chi(L^\omega)$.

Proposition 4 *Let L^ω and M^ω be two rational adherences. The following assertions are equivalent:*

- (i) $L^\omega = M^\omega$
- (ii) $Per_\chi(L^\omega) = Per_\chi(M^\omega)$

Proof (i) \Rightarrow (ii) If $L^\omega = M^\omega$, we obtain immediately $Per_\chi(L^\omega) = Per_\chi(M^\omega)$. Conversely, if $Per_\chi(L^\omega) = Per_\chi(M^\omega)$, from corollary 3, we deduce that $Pref(L^\omega) = Pref(Per_\chi(L^\omega)) = Pref(Per_\chi(M^\omega)) = Pref(M^\omega)$. Moreover, we know that two rational adherences are equal if and only if their sets of prefixes are equal (corollary 1). \square

Example Let $K = a^*ba^\omega + a^\omega$ and $L = a^\omega$ be two adherences. K is clearly distinct from L . However, $Per_\chi(K) = Per_\chi(L) = a^+$. It is allowed since K is not an ω -power.

4.2 Rational deterministic ω -languages

By considering rational deterministic ω -languages, we obtain complementary results. They are recognized by deterministic Büchi automata and form a proper subclass of the class of rational ω -languages [4]. Note that a rational adherence L^ω is a deterministic ω -power. A fortiori, a finitely ω -generated ω -language is a deterministic rational ω -power.

Proposition 5 [6] *If L^ω is a rational deterministic ω -language, then every word u from $\chi(L^\omega)$ belongs to some ω -generator G of L^ω .*

Moreover, as mentionned in [6], in the case of a non deterministic rational ω -power, it may occur that L^ω has a greatest ω -generator different from $\chi(L^\omega)$.

Recall that in the general case, it is proved in [7] that a rational ω -language L^ω admits a finite number of maximal ω -generators, with respect to inclusion.

Corollary 4 *Let L^ω be a deterministic rational ω -language and let M_1, \dots, M_n denote its maximal ω -generators, then:*

$$\chi(L^\omega) = \bigcup_{i=1}^n M_i$$

Proof According to proposition 5, for every $u \in \chi(L^\omega)$, there exists an ω -generator G of L^ω such that $u \in G$: $\chi(L^\omega) \subseteq \bigcup_{i=1}^n M_i$. Conversely, the set obtained as the union of all the ω -generators of L^ω is included in the upper bound $\chi(L^\omega)$. \square

If $\chi(L^\omega)$ is not a semigroup, we can remark that the elements u, v in $\chi(L^\omega)$ verifying $uv \notin \chi(L^\omega)$ do not belong to the intersection of maximal ω -generators. Indeed, for every $u, v \in \chi(L^\omega)$ with $uv \notin \chi(L^\omega)$, thus $(uv)^\omega$ does not belong to L^ω . The two words u and v cannot be in the same ω -generator and so not in the same maximal one. So they do not belong to the intersection of these maximal elements.

Proposition 6 *Let L^ω be a rational deterministic ω -language. There exists an integer n such that every ω -generator G of L^ω verifies:*

$$\forall m \geq n, \quad G^m \subseteq I$$

where I denotes the intersection of the maximal ω -generators of L^ω .

Proof Let $\mathcal{A} = (\Sigma, Q, \delta, q_0, T)$ be a deterministic Büchi automaton which recognizes L^ω . Since \mathcal{A} is deterministic, its transition relation is a function from $Q \times \Sigma$ into Q . It associates to (q, σ) the element $\delta(q, \sigma)$, and it can be prolongable to Σ^* . We set $n = \text{card}(Q) + 1$. We intend to prove that, for each ω -generator G of L^ω , $G^n \subseteq I$. We write $K = \{q, q = \delta(q_0, u), u \in \text{Stab}(L^\omega)\}$. We note $l(q, v)$ the set of states which occur in a run from the state q and labelled by v in \mathcal{A} . We start with the proof of the following fact, close to a lemma which can be found in [6]:

Lemma 3 $\forall q \in K, \quad \forall v \in G^n, \quad l(q, v) \cap T \neq \emptyset.$




Figure 1: Illustration for the proof of lemma 3

Proof of the lemma: Consider $q \in K$ and $v = v_1 \dots v_n$ with $\forall i, 1 \leq i \leq n, v_i \in G$. As $\text{Card}(\{\delta(q, v_1 \dots v_i), 1 \leq i \leq n\}) < n$, so there exist two different integers j and j' , $j < j'$ such that $\delta(q, v_1 \dots v_j) = \delta(q, v_1 \dots v_{j'})$. Moreover, $v_1 \dots v_j (v_{j+1} \dots v_{j'})^\omega \in G^\omega = L^\omega$, then, $\inf(l(q, v_1 \dots v_j (v_{j+1} \dots v_{j'})^\omega)) \cap T \neq \emptyset$. By the determinism of \mathcal{A} , we deduce that: $l(q, v_1 \dots v_j, v_{j+1} \dots v_{j'}) \cap T \neq \emptyset$. At last, $l(q, v_1 \dots v_n) \cap T \neq \emptyset$ which achieves the proof of the lemma.

Now, we prove that $G^n \subseteq I$ (the same proof holds for every $m > n$). Assume that there exists an element x in $G^n \setminus I$. Then, there exists a maximal ω -generator M of L^ω such that $x \in G^n \setminus M$. We want to prove $(M + x)^\omega = L^\omega$.

It is clear that $G^\omega \subseteq (M + x)^\omega$. Conversely, $w \in (M + x)^\omega$ with $(w_i)_{i>0}$ a factorization of w over $M + x$. Two different cases may occur:

- if $\text{Card}(\{w_i, w_i = x\}) = \infty$. $w \in (Mx)^\omega$ then $w = m_1 x m_2 x \dots$, $m_i \in M$. Moreover, $\forall i > 0 \delta(q_0, m_1 x \dots m_i) \in K$ and so, according to the previous lemma, if $\forall q \in K, \forall i > 0 l(\delta(q, m_1 x \dots m_i), x) \cap T \neq \emptyset$. Consequently, $w \in L^\omega$.
- if $\text{Card}(\{w_i, w_i = x\})$ is finite, then $w \in (M + x)^* M^\omega \subseteq L^\omega$.

We obtain $(M + x)^\omega = L^\omega$, which contradicts the maximality of M . \square

Theorem 2 *Let L^ω be a rational deterministic ω -language. The intersection I of maximal ω -generators is still an ω -generator of L^ω .*

Proof For every $j \in \{1, \dots, n\}$, $(\bigcap_{i=1}^n M_i) = I \subseteq M_j$ then $I^\omega \subseteq M_j^\omega = L^\omega$. Conversely and using the previous lemma, there exists an integer n such that $L^n \subseteq I$ that is $L^\omega \subseteq I^\omega$. \square

References

- [1] A. Arnold. A syntactic congruence for rational ω -languages. *Theoretical Computer Science*, 39:333–335, 1985. Note.

- [2] J.R. Büchi. On a decision method in restricted second order arithmetics. In *International Congress on Logic, Methodology and Philosophy of Science*, pages 1–11. Stanford University Press, 1960.
- [3] J. Devolder-Muchembled. *Codes, mots infinis et bi-infinis*. PhD thesis, Université de Lille, 1993.
- [4] S. Eilenberg. *Automata, Languages and Machines*, volume A. Academic Press, 1974.
- [5] M. Latteux and E. Timmerman. Finitely generated ω -langages. *Information Processing Letters*, 23:171–175, 1986.
- [6] I. Litovsky. *Générateurs des langages rationnels de mots infinis*. PhD thesis, Université de Lille, 1988.
- [7] I. Litovsky and E. Timmerman. On generators of rational ω -power languages. *Theoretical Computer Science*, 53:187–200, 1987.
- [8] R. Mac Naughton. Testing and generating infinite sequences by a finite automaton. *Information and Control*, 9:521–530, 1966.
- [9] M. Nivat. Sur les ensembles de mots infinis engendrés par une grammaire algébrique. *R.A.I.R.O. Informatique Théorique*, 12:259–278, 1978.